

Updated GLM standardised CPUE abundance indices for orange roughy for known aggregations off Namibia from 1994 to 2006

Anabela Brandão and Doug S. Butterworth

*Marine Resource Assessment & Management Group (MARAM)
Department of Mathematics and Applied Mathematics
University of Cape Town
Rondebosch, 7701, Cape Town*

February 2007

Abstract

GLM analyses are used to standardise the CPUE data for Namibian orange roughy in a manner that deals with tows that record zero catch of orange roughy. The possibility of there being a “learning” period of lower CPUE for a new vessel when it enters the fishery is taken into account. Further, to allow for areal expansion of the fishery at each aggregation, sub-aggregations are defined and CPUE trends estimated separately for each. The “zero” method for combining the results for the various sub-aggregations to provide a single index for an aggregation is applied. The standardised CPUE value for 2005 (i.e. the July 2004 – June 2005 fishing year) for *Johnies* is the highest it has been in the last six years. The 2006 index is down from the previous year. For *Frankies* the 2006 index is up from that for 2004 (no 2005 index is available). No indices for *Hotspot* and *Rix* are available for the last two years.

Introduction

A delta-lognormal model, as first proposed in Brandão and Butterworth (2002), is used to standardise the commercial orange roughy CPUE data. This type of model addressed two problems encountered in the analyses for this fishery: i) a considerable number of tows with

zero catches and ii) the areal distribution of effort shifting within and even beyond previously defined aggregations (especially notable for the *Johnies* aggregation). These standardised CPUE indices of abundance are then used as an input to a population model to assess the state of the stock (Brandão and Butterworth 2007). In this paper, the results of the updated standardised CPUE indices for orange roughy taking additional data for the 2005 and 2006 fishing season into account are presented.

The Model

The model applied to the CPUE time series of data for Namibian orange roughy is a delta-lognormal which takes into account the presence of tows with zero catch as described by Lo *et al.* (1992) and Stone and Porter (1999).

The delta distribution is often used in instances when there are a considerable number of zero observations, for which zero and non-zero data are consequently treated separately. Final estimates of abundance are obtained from the product of the proportion and the mean of non-zero observations. For the delta-lognormal model, two linear models are fitted to the commercial CPUE data, one to estimate the proportion of tows for which there is a positive catch, and the other to estimate the standardised CPUE for orange roughy for tows that have a positive catch.

Relative abundance indices of orange roughy are then given by:

$$CPUE_y = \sum_{agg} (CPUE_{agg,y}^{+ve} Prop_{agg,y}^{non-zero}) A_{agg} \quad (1)$$

where:

$CPUE_{agg,y}^{+ve}$ is the standardised CPUE index for tows which have positive catches for a given sub-aggregation,

$Prop_{agg,y}^{non-zero}$ is the standardised measure of the proportion of tows that have positive catches for a given sub-aggregation, and

A_{agg} is the geographical area for a given sub-aggregation (Table 1).

Standardised indices for the component related to the CPUE of positive catches were obtained by fitting a lognormal model that allows for possible differences in abundance trends in orange roughy in the various aggregations, and assume the possibility that vessels might operate differently in their first year in the fishery, but have the same degree of “effectiveness” in all subsequent years. Brandão and Butterworth (2003) found that only the

vessel *Whitby* showed a significant difference in its first year of operation. New vessels that have operated in the fishery since this analysis have not shown a significant difference in their first year of operation and therefore only the *Whitby* vessel is differentiated with respect to its first year in the fishery and all subsequent years. The model to estimate the standardised index of positive catches is thus given by:

$$\ln(\text{CPUE}^{+ve}) = \mu + \alpha_{\text{vessel}} + \beta_y + \gamma_{\text{month}} + \lambda_{\text{agg}} + \eta_{y \times \text{agg}} + \varepsilon \quad (2)$$

where:

μ is the intercept,

vessel is a factor with 14 levels associated with each of the vessels that have operated in the fishery:

Bell Ocean II

Conbaroya Quarto

Concasa

Dantago

Emanguluko

Harvest Nicola

Hurinis

Petersen

Sea Flower

Southern Aquarius

Ulzama

Whitby (first year)

Whitby (subsequent years)

Will Watch,

y is a factor with 13 levels associated with the “fishing years” 1994–2006 (note: “1996”, for example, refers to the period July 1996 to June 1997),

month is a factor with 12 levels (January– December),

agg is a factor with 12 levels associated with the four aggregations and their sub-aggregations:

Johnies: Johnies1

Johnies2

Johnies3

Johnies4

Frankies: 21 Jump Street

Frankies Flats

Frankies Outer

Three Sisters

Smifton

Rix: Rix Inner

Rix Outer

Hotspot,

$y \times agg$ is the interaction between year and aggregation (this allows for the possibility of different temporal trends for the different sub-aggregations), and

ε is an error term assumed to be normally distributed.

In the case of the orange roughy tow data, the proportion of tows with a positive catch is either “0” or “1” for an individual tow, and therefore a model for the proportion positive assuming binomially distributed errors is considered, given by:

$$\ln\left(\frac{Prop^{non-zero}}{1 - Prop^{non-zero}}\right) = \mu + \alpha_{vessel} + \beta_y + \gamma_{month} + \lambda_{agg} + \eta_{y \times agg} + \zeta \quad (3)$$

where

ζ is an error term assumed to be binomially distributed.

Standardised measures of the abundance of orange roughy in positive tows for a given (sub)-aggregation are estimated by calculating:

$$C\hat{P}UE_{agg,y}^{+ve} = \exp\left[\hat{\mu} + \hat{\beta}_y + \hat{\lambda}_{agg} + \hat{\eta}_{y \times agg}\right] \psi_y^{+ve} \quad (4)$$

where in this application standardisation is with respect to the vessel *Southern Aquarius* and to the month of *August*, and where

ψ_v^{+ve} is a correction factor for bias (Lo *et al.* 1992), given by:

$$\psi_y^{+ve} = g_m \left[\frac{m+1}{2m} \left(\hat{\xi}^2 - \hat{\xi}_{\hat{\theta}}^2 \right) \right] \quad (5)$$

where

$\hat{\xi}^2$ is the residual variance,

m is the degrees of freedom for the estimate of residual variance,

$\hat{\theta}$ is given by $\hat{\mu} + \hat{\beta}_y + \hat{\lambda}_{agg} + \hat{\eta}_{y \times agg}$,

ξ_{θ}^2 is the variance of $\hat{\theta}$, and

$$g_m(t) = \sum_{p=0}^{\infty} \left[\frac{m^p (m+2p)}{m(m+2) \cdots (m+2p)} \left(\frac{m}{m+1} \right)^p \frac{t^p}{p!} \right]$$

where t is the argument of the function.

Standardised measures of the proportion of positive catches of orange roughy are given by:

$$\hat{Prop}_{agg,y}^{non-zero} = \frac{\exp[\hat{\mu} + \hat{\beta}_y + \hat{\lambda}_{agg} + \hat{\eta}_{y \times agg}]}{1 + \exp[\hat{\mu} + \hat{\beta}_y + \hat{\lambda}_{agg} + \hat{\eta}_{y \times agg}]} \quad (6)$$

Model Implementation

To take into account movement of orange roughy within a known aggregation, the analyses in Brandão and Butterworth (2002) considered not only tows that lie within the inner strata of an aggregation, but also tows that take place in the outer strata of the aggregation. The levels of the factor for aggregations in the GLMs thus correspond to the various sub-aggregations. The definitions of aggregations and their sub-aggregations given by Brandão and Butterworth (2002) are used in this paper.

Commercial tow information inside the known aggregations of orange roughy in Namibia for the fishing years (July–June) 1994 to 2006, as provided by NatMIRC, has been used. As a restriction is applied to the data records used in the GLM analyses that there must be 20 or more records within a fishing year in each sub-aggregation, insufficient data were available for the analyses to include the 2005 fishing year for *Frankies* and *Hotspot*, as well as 2006 for *Hotspot*. A total of 18 201 tows was available for the analyses. Of these, 15 348 recorded a non-zero catch. Bottom distances were calculated from the GPS positions for each tow. For tows that did not have haul positions (the majority of tows in the last few years), but did have bottom time information, bottom distances were calculated by the following regression relationship:

$$\text{Bottom distance [km]} = \text{bottom time [h]} * 5.6082 + 0.1259$$

developed in earlier analyses (Brandão and Butterworth 2003).

GLM Results and Discussion

The lognormal model applied to tows with a positive catch (equation (2)) accounts for 46.5% of the total variation of orange roughy positive CPUE. Table 2 shows the parameter estimates obtained for the factor vessel for the CPUE of positive catches and for the proportion of positive tows. Table 3 shows the index of abundance provided by the delta-lognormal model assuming binomial errors for the proportion positive for each aggregation. Observations are not available for all years in all of the sub-aggregations. In this paper only the “zero” method (Brandão and Butterworth, 2002) of combining the standardised CPUE indices from each individual sub-aggregation to obtain a standardised CPUE index for each aggregation was used to deal with such empty cells (i.e. assume that empty cells mean that there was no orange roughy in those areas for those years). The overall standardised index for each aggregation is obtained by summing the standardised CPUE for each sub-aggregation multiplied by its associated geographical area (equation(1)).

The restriction applied to the CPUE data used in the GLM analyses implies that no CPUE records were available for the *Hotspot* aggregation for the years 2005 to 2006. Table 3 also shows the nominal CPUE series for Hotspot, including the tows that were performed in 2005 and 2006.

Figures 1 to 4 show the index of abundance provided by the delta-lognormal model assuming binomial errors for the proportion positive for each aggregation. For comparison purposes, nominal CPUE series are also shown in Figures 1 to 4. Differences in the series are most marked in the first few years of the series. Figure 4b shows the nominal CPUE series for Hotspot that includes data for 2005 and 2006 (with the vertical axis scale changed so that the CPUE trends in the later years becomes clearer).

The standardised CPUE value for 2005 for *Johnies* is the highest of the last six years, but the 2006 index is lower and in the range of those for 2001 to 2003. However, the 2006 fishing year is incomplete (ending only in June 2007) and therefore this value might change as further data become available. The standardised CPUE values for 2006 for *Frankies* is higher than that for 2004 (no index is available for 2005 because of minimum data requirements applied in the GLM analyses). No indices are available for *Hotspot* for the years 2005 and 2006 for these same minimum data requirement reasons. However, nominal CPUE indices for these years show that the 2005 value is near to that for 1997, and the

2006 value is (2005 apart) the highest since 1998. No data are available for *Rix* since 2004 as this aggregation was closed for commercial fishing on 1st August 2004.

Acknowledgements

Data were provided by NatMIRC for this study. Funding from the Namibian Deepwater Fishing Industry is gratefully acknowledged.

References

- Brandão, A. and Butterworth, D.S. 2002. Standardised CPUE abundance indices for orange roughy off Namibia based on lognormal and delta-lognormal models. Namibian Ministry of Fisheries and Marine Resources document: DWFWG/WkShop/Feb02/Doc 1.
- Brandão, A. and Butterworth, D.S. 2003. Application of lognormal and delta-lognormal linear models to standardised CPUE abundance indices (1994 to 2002) for orange roughy off Namibia. Namibian Ministry of Fisheries and Marine Resources document: DWFWG/WkShop/Mar03/Doc 2.
- Brandão, A. and Butterworth, D.S. 2007. Age-structured production model assessments and projections including updated parameters to model the intermittent aggregation of Namibian orange roughy. Namibian Ministry of Fisheries and Marine Resources document: DWFWG/WkShop/Feb07/Doc 5.
- Lo, N.C., Jacobson, L.D., and Squire, J.L. 1992. Indices of relative abundance from fish spotter data based on delta-lognormal models. *Can. J. Fish. Aquat. Sci.* 49: 2515–2526.
- Stone, H.H. and Porter, J.M. (1999) Standardized CPUE indices for Canadian bluefin tuna fisheries based on commercial catch rates. *International Commission for the Conservation of Atlantic Tunas, collective Volume of Scientific Papers: XLIX (2): 206–221.*

Table 1. Geographical area for each sub-aggregation of orange roughy off Namibia.

Aggregation	Sub-aggregation	Area (km²)
<i>Johnnies</i>	<i>Johnnies1</i>	82.8
	<i>Johnnies2</i>	457.2
	<i>Johnnies3</i>	198.2
	<i>Johnnies4</i>	587.1
<i>Frankies</i>	<i>21 Jump Street</i>	39.2
	<i>Frankies Flats</i>	17.8
	<i>Frankies Outer</i>	1 255.0
	<i>Three Sisters</i>	39.6
	<i>Smifton</i>	15.8
<i>Rix</i>	<i>Rix Inner</i>	99.4
	<i>Rix Outer</i>	685.6
<i>Hotspot</i>	<i>Hotspot Inner</i>	97.3
	<i>Hotspot Outer*</i>	89.0

* Too few tows fall within the *Hotspot Outer* sub-aggregation for specific account to be taken of this sub-aggregation, and therefore these tows are omitted from the GLM analyses.

Table 2. Parameter estimates for the vessel factor when the lognormal model is applied to tows with a positive catch (equation (2)) and the model for the proportion positive (equation (3)) are fitted.

Vessel	Vessel factor = $e^{\alpha_{vessel}}$ (positive catches)	Vessel factor = $e^{\alpha_{vessel}}$ (proportion positive)
<i>Bell Ocean II</i>	0.382	0.173
<i>Conbaroya Cuarto</i>	0.293	1.228
<i>Concasa</i>	0.186	1.350
<i>Dantago</i>	0.311	0.776
<i>Emanguluko</i>	0.447	1.129
<i>Harvest Nicola</i>	0.212	0.494
<i>Hurinis</i>	0.318	0.627
<i>Petersen</i>	0.427	4.293
<i>Sea Flower</i>	0.502	2344*
<i>Southern Aquarius</i>	1.000	1.000
<i>Ulzama</i>	1.240	0.501
<i>Whitby</i> (first year)	0.995	1.029
<i>Whitby</i> (subsequent years)	0.333	172*
<i>Will Watch</i>	0.995	2038*

* Note: These large values are not unrealistic, but rather are a consequence of the logit transformation used in equation (3) [which restricts the final factor applied to lie between 0 and 1] and the fact that these three vessels had no records of zero tows.

Table 3. Standardised CPUE series (normalised to their mean over the years considered) for the *Johnnies*, *Frankies*, *Rix* and *Hotspot* aggregations obtained by fitting the delta-lognormal model, assuming binomial errors for the proportion positive, to the observed CPUE data for Namibian orange roughy. The “zero” method for dealing with years in which no observations were made in the sub-aggregations is considered. The *Frankies* aggregation was closed in 1999 and has been partially reopened since 2002 and fully reopened since 2005 (calendar years). Therefore the indices for the fishing years that span those calendar years are based on very few data. For comparison, the nominal CPUE series for *Hotspot* (that includes data for 2005 and 2006 that had been excluded in the GLM analysis) is also shown.

Year	<i>Johnnies</i>	<i>Frankies</i>	<i>Rix</i>	<i>Hotspot</i>	<i>Hotspot (nominal)</i>
1994	6.411	—		6.236	6.038
1995	1.006	1.354	0.518	1.815	2.067
1996	1.382	4.797	0.676	0.941	1.094
1997	1.827	1.499	4.415	0.333	0.584
1998	0.662	0.715	1.914	0.524	0.314
1999	0.296	0.325	0.379	0.277	0.208
2000	0.256	—	0.393	0.110	0.089
2001	0.142	0.474	0.280	0.178	0.162
2002	0.179	0.167	0.282	0.378	0.201
2003	0.151	0.474	0.144	0.099	0.089
2004	0.067	0.024	—†	0.109	0.155
2005	0.456	—*	—†	—*	0.520
2006	0.166	0.171	—†	—*	0.242

* There were too few tows to be included in the GLM analyses, thus no index is available for this year.

† The Rix aggregation has been closed to commercial fishing since 1st August 2004 (calendar year).

Johnnies




Figure 1. Index of abundance for the *Johnnies* aggregation (normalised to its mean over the thirteen year period) for Namibian orange roughly obtained from fitting the delta-lognormal model assuming binomial errors for the proportion positive. Results are shown for the “zero” method of dealing with empty cells when combining the indices from sub-aggregations. For comparison, the nominal CPUE series is also shown.

Frankies




Figure 2. Index of abundance for the *Frankies* aggregation (normalised to its mean over the twelve year period) for Namibian orange roughly obtained from fitting the delta-lognormal model assuming binomial errors for the proportion positive. Results are shown for the “zero” method of dealing with empty cells when combining the indices from sub-aggregations. For comparison, the nominal CPUE series obtained is also shown.




Figure 3. Index of abundance for the *Rix* aggregation (normalised to its mean over the twelve year period) for Namibian orange roughly obtained from fitting the delta-lognormal model assuming binomial errors for the proportion positive. Results are shown for the “zero” method of dealing with empty cells when combining the indices from sub-aggregations. For comparison, the nominal CPUE series is also shown.




Figure 4a. Index of abundance for the *Hotspot* aggregation (normalised to its mean over the thirteen year period) for Namibian orange roughly obtained from fitting the delta-lognormal model assuming binomial errors for the proportion positive. For comparison, the nominal CPUE series is also shown.




Figure 4b. Index of abundance for the *Hotspot* aggregation (normalised to its mean over the thirteen year period) for Namibian orange roughly obtained from fitting the delta-lognormal model assuming binomial errors for the proportion positive. For comparison, the nominal CPUE series (that includes data for 2005 and 2006 that had been excluded in the GLM analysis) is also shown. For clarification of the indices in the later years, the vertical axis scale has been modified.