

Quantifying chaos using Lagrangian descriptors

Haris Skokos

**Department of Mathematics and Applied Mathematics
University of Cape Town
Cape Town, South Africa**

E-mail: haris.skokos@uct.ac.za

URL: http://math_research.uct.ac.za/~hskokos/

Work in collaboration with:

Sebastian Zimmer, Malcolm Hillebrand, Arnold Ngapasare (University of Cape Town, South Africa)

Matthaios Katsanikas (Academy of Athens, Greece)

Stephen Wiggins (University of Bristol, UK & United States Naval Academy, USA)

**4th International Conference on Integrable Systems and Nonlinear Dynamics (ISND) 2023
27 September 2023, Yaroslavl, Russia**

University of Cape Town (UCT)



University of Cape Town (UCT)



Cape Town



Outline

- **Lagrangian descriptors (LDs)**
- **Smaller Alignment Index (SALI)**
- **Chaos diagnostics based on LDs:**
 - ✓ the difference of LDs of neighboring orbits
 - ✓ the ratio of LDs of neighboring orbits
- **Application to the Hénon – Heiles system**
- **Summary**

Lagrangian descriptors (LDs)

The computation of LDs is based on the accumulation of some positive scalar value along the path of individual orbits.

Consider an N dimensional continuous time dynamical system

$$\dot{x} = \frac{dx(t)}{dt} = f(x, t)$$

The Arclength Definition [Madrid & Mancho, Chaos (2009) – Mendoza & Mancho, PRL (2010) – Mancho et al., Commun. Nonlin. Sci. Num. Simul. (2013)].

Forward time LD:

$$LD^f(x, \tau) = \int_0^\tau \|\dot{x}(t)\| dt$$

Backward time LD:

$$LD^b(x, \tau) = \int_{-\tau}^0 \|\dot{x}(t)\| dt$$


Combined LD:

$$LD(x, \tau) = LD^b(x, \tau) + LD^f(x, \tau)$$

LDs: 1 degree of freedom (dof) Hamiltonian

$$H(q, p) = \frac{1}{2} (p^2 - q^2)$$



The system has a hyperbolic fixed point at the origin. The LDs can be used to display the stable and unstable manifolds of this point.



LDs: 1 dof Duffing Oscillator

$$H(x, y) = \frac{1}{2}y^2 + \frac{1}{4}x^4 - \frac{1}{2}x^2$$

The system has three equilibrium points: a saddle located at the origin and two diametrically opposed centers at the points $(\pm 1, 0)$.



From Agaoglou et al. 'Lagrangian descriptors: Discovery and quantification of phase space structure and transport', 2020, <https://doi.org/10.5281/zenodo.3958985>

The **location of the stable and unstable manifolds** can be extracted from the ridges of the **gradient field of the LDs** since they are located at points where the forward and the backward components of the LD are non-differentiable.

Lagrangian descriptors (LDs)


The ‘ p -norm’ Definition [Lopesino et al., Commun. Nonlin. Sci. Num. Simul. (2015) – Lopesino et al., Int. J. Bifurc. Chaos (2017)].

Combined LD (usually $p=1/2$):

$$LD(x, \tau) = \int_{-\tau}^{\tau} \left(\sum_{i=1}^N |f_i(x, t)|^p \right) dt$$

Hénon-Heiles system: $H = \frac{1}{2}(p_x^2 + p_y^2) + \frac{1}{2}(x^2 + y^2) + x^2 y - \frac{1}{3}y^3$

Stable and **unstable** manifolds for $H=1/3, \tau=10$.



The Smaller Alignment Index (SALI)

Consider the **2N-dimensional** phase space of a conservative dynamical system (**symplectic map or Hamiltonian flow**).

An orbit in that space with initial condition :

$$P(0)=(x_1(0), x_2(0), \dots, x_{2N}(0))$$

and **a deviation vector**

$$v(0)=(\delta x_1(0), \delta x_2(0), \dots, \delta x_{2N}(0))$$

The evolution in time (in maps the time is discrete and is equal to the number n of the iterations) of **a deviation vector** is defined by:

- the **variational equations** (for Hamiltonian flows) and
- the equations of the **tangent map** (for mappings)

Definition of the SALI

We follow the evolution in time of two different initial deviation vectors ($\mathbf{v}_1(0)$, $\mathbf{v}_2(0)$), and define SALI [S., J. Phys. A (2001) – S & Manos, Lect. Notes Phys. (2016)] as:

$$\text{SALI}(t) = \min\{\|\hat{\mathbf{v}}_1(t) + \hat{\mathbf{v}}_2(t)\|, \|\hat{\mathbf{v}}_1(t) - \hat{\mathbf{v}}_2(t)\|\}$$

where

$$\hat{\mathbf{v}}_1(t) = \frac{\mathbf{v}_1(t)}{\|\mathbf{v}_1(t)\|}$$

When the two vectors become collinear

$$\text{SALI}(t) \rightarrow 0$$

SALI – Hénon-Heiles system

As an example, we consider the 2D Hénon-Heiles system:



$$H = \frac{1}{2}(p_x^2 + p_y^2) + \frac{1}{2}(x^2 + y^2) + x^2y - \frac{1}{3}y^3$$

For $E=1/8$ we consider the orbits with initial conditions:


Regular orbit, $x=0$, $y=0.55$, $p_x=0.2417$, $p_y=0$

Chaotic orbit, $x=0$, $y=-0.016$, $p_x=0.49974$, $p_y=0$

Chaotic orbit, $x=0$, $y=-0.01344$, $p_x=0.49982$, $p_y=0$



SALI – Hénon-Heiles system



Behavior of the SALI

2D maps

SALI $\rightarrow 0$ both for regular and chaotic orbits

following, however, completely different time rates which allows us to distinguish between the two cases.

Hamiltonian flows and multidimensional maps

SALI $\rightarrow 0$ for chaotic orbits

SALI $\rightarrow \text{constant} \neq 0$ for regular orbits

Using LDs to quantify chaos

We consider orbits on a finite **grid of an $n(\geq 1)$ -dimensional subspace** of the **$N(\geq n)$ -dimensional phase space** of a dynamical system and their LDs.

Any non-boundary point x in this subspace has **$2n$ nearest neighbors**

$$y_i^{\pm} = x \pm \sigma^{(i)} e^{(i)}, \quad i = 1, 2, \dots, n,$$

where $e^{(i)}$ is the i th usual basis vector in \mathbb{R}^n and $\sigma^{(i)}$ is the distance between successive grid points in this direction.


The **difference D_L^n** of neighboring orbits' LDs:

$$D_L^n(x) = \frac{1}{2n} \sum_{i=1}^n \frac{|LD^f(x) - LD^f(y_i^+)| + |LD^f(x) - LD^f(y_i^-)|}{LD^f(x)}.$$

The **ratio R_L^n** of neighboring orbits' LDs:



$$R_L^n(x) = \left| 1 - \frac{1}{2n} \sum_{i=1}^n \frac{LD^f(y_i^+) + LD^f(y_i^-)}{LD^f(x)} \right|.$$

Application: Hénon-Heiles system







Application: Hénon-Heiles system

$H=1/8$




Variation of LDs with regard to initial conditions.
regular regions: smooth
chaotic regions: erratic
 [also see Montes et al., Commun. Nonlin. Sci. Num. Simul. (2021)]

LDs for $\tau=10^3$




SALI for $\tau=10^6$
(inset $\tau=10^3$)

Application: Hénon-Heiles system




Misclassified orbits (< 10%)



Application: Hénon-Heiles system

A quantity related to **the second derivative of the LDs** was introduced in Daquin et al., Physica D (2022) and was used in Hillebrand et al., Chaos (2022) and Zimper et al., Physica D (2023):

$$\|\Delta LD\|(x) = \left| \frac{LD^f(y_i^+) - 2LD^f(x) + LD^f(y_i^-)}{\sigma^2} \right|.$$



Summary

- ✓ We introduced and successfully implemented computationally efficient ways to **effectively identify chaos** in conservative dynamical systems **from the values of LDs at neighboring initial conditions**.
- ✓ From the distributions of the indices' values we determine appropriate **threshold values**, which allow the characterization of orbits as regular or chaotic.
- ✓ Both the indices **faced problems** in correctly revealing the nature of some orbits mainly **at the borders of stability islands**.
- ✓ Both indices show **overall very good performance**, as their classifications are in accordance with the ones obtained by **the SALI at a level of at least 90% agreement**.
- ✓ **Advantages:**
 - **Easy to compute** (actually only the forward LDs are needed).
 - **No need to know and to integrate the variational equations**.
- ✓ These methods has also been successfully applied to **2D and 4D symplectic maps** [Hillebrand et al., Chaos (2022) – Zimmer et al., Physica D (2023)]

References

- S., J. Phys. A, 34, 10029 (2001)
- Madrid & Mancho, Chaos, 19, 013111 (2009)
- Mendoza & Mancho, PRL, 105, 038501 (2010)
- Mancho, Wiggins, Curbelo & Mendoza, Commun. Nonlin. Sci. Num. Simul., 18, 3530 (2013)
- Lopesino, Balibrea, Wiggins & Mancho, Commun. Nonlin. Sci. Num. Simul., 27, 40 (2015)
- S. & Manos, Lect. Notes Phys., 915, 129 (2016)
- Lopesino, Balibrea-Iniesta, García-Garrido, Wiggins & Mancho, Int. J. Bifurc. Chaos, 27, 1730001 (2017)
- Agaoglou, Aguilar-Sanjuan, García-Garrido, González-Montoya, Katsanikas, Krajňák, Naik & Wiggins, ‘Lagrangian descriptors: Discovery and quantification of phase space structure and transport’, <https://doi.org/10.5281/zenodo.3958985> (2020)
- Montes, Revuelta & Borondo, Commun. Nonlin. Sci. Num. Simul., 102, 105860 (2021)
- Daquin, Pédenon-Orlanducci, Agaoglou, García-Sánchez & Mancho, Physica D, 442, 133520 (2022)

Hillebrand, Zimper, Ngapasare, Katsanikas, Wiggins & S.: Chaos, 32, 123122 (2022),
‘Quantifying chaos using Lagrangian descriptors’

Zimper, Ngapasare, Hillebrand, Katsanikas, Wiggins & S.: Physica D, 453, 133833 (2023),
‘Performance of chaos diagnostics based on Lagrangian descriptors. Application to the 4D standard map’